



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

WEST INDIES.

Report from Castries, St. Lucia Island—Sanitary conditions good.

Acting Assistant Surgeon Maylie reports, June 17:

Sanitary condition of port very good; no quarantinable diseases.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

AFRICA—*Zanzibar*.—Ten days ended May 31, 1907. Estimated population, 75,000. Total number of deaths, 31, including 3 from tuberculosis.

AUSTRALIA—*Freemantle and Perth*.—Month of January, 1907. Estimated population, 81,700. Total number of deaths, 119, including diphtheria 4 and 4 from enteric fever. Month of February, 1907. Total number of deaths, 97, including diphtheria 2, enteric fever 6, measles 1, and 2 from whooping cough. Month of March, 1907. Total number of deaths, 86, including diphtheria 2, enteric fever 5, measles 1, and 1 from whooping cough.

CENTRAL AMERICA—*Salvador—San Salvador*.—Three months ended December 31, 1906. Estimated population, 58,753. Total number of deaths, 370, including whooping cough 2 and 38 from tuberculosis. Three months ended March 31, 1907. Total number of deaths, 389, including 29 from tuberculosis.

COSTA RICA—*San Jose*.—Month of May, 1907. Estimated population, 25,525. Total number of deaths, 64, including enteric fever 1, and 3 from tuberculosis.

FRANCE—*Cannes*.—Month of May, 1907. Estimated population, 37,361. Total number of deaths not reported. Two deaths from smallpox reported.

Nice.—Month of May, 1907. Estimated population, 150,881. Total number of deaths, 223, including enteric fever 2, smallpox 1, and 35 from tuberculosis.

GIBRALTAR.—Two weeks ended June 23, 1907. Estimated population, 27,385. Total number of deaths 10. No deaths from contagious diseases.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended June 15, 1907, correspond to an annual rate of 13.1 per 1,000 of population, which is estimated at 16,024,458.

London.—One thousand one hundred and seventy-two deaths were registered during the week, including measles 56, scarlet fever 13, enteric fever 2, diphtheria 8, whooping cough 53, tuberculosis 157, and 19 from diarrhea. The deaths from all causes correspond to an annual rate of 12.8 per 1,000. In Greater London 1,630 deaths were regis-

tered. In the "outer ring" the deaths included 4 from measles, 2 from scarlet fever, 4 from diphtheria, and 5 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended June 15, 1907, in the 21 principal town districts of Ireland, was 21.2 per 1,000 of the population, which is estimated at 1,117,547. The lowest rate was recorded in Galway, viz, 3.9, and the highest in Sligo, viz, 62.4 per 1,000. In Dublin and suburbs 191 deaths were registered, including measles 3, whooping cough 8, and 38 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended June 15, 1907, correspond to an annual rate of 17.2 per 1,000 of the population, which is estimated at 1,812,171. The highest rate of mortality was recorded in Dundee, viz, 22.0, and the lowest in Greenock, viz, 11.0 per 1,000. The aggregate number of deaths registered from all causes was 599, including diphtheria 3, enteric fever 1, measles 5, and 43 from whooping cough.

JAPAN—Formosa.—Two weeks ended June 1, 1907. Estimated population, 3,050,004. Total number of deaths not reported. Three deaths from enteric fever and 299 from plague reported.

MALTA.—Two weeks ended June 15, 1907. Estimated population, 206,690. Total number of deaths, 172. No deaths from contagious diseases reported.

MAURITIUS.—Month of March, 1907. Estimated population, 375,400. Total number of deaths, 1,058, including enteric fever 3, plague 3, and 2 from whooping cough.

SPAIN—Huelva.—Month of May, 1907. Estimated population, 24,000. Total number of deaths, 67, including enteric fever 2, measles 2, smallpox 1, and 11 from tuberculosis.

Seville.—Month of May, 1907. Estimated population, 148,315. Total number of deaths, 469, including diphtheria 5, enteric fever 5, measles 3, whooping cough 6, smallpox 15, and 72 from tuberculosis.

SWITZERLAND.—Week ended June 8, 1907. Reports from 18 cities. having an aggregate population of 867,056, show as follows: Total number of deaths, 230, including diphtheria 4, enteric fever 1, measles 5, and 42 from tuberculosis.

WEST INDIES—Curaçao.—Two weeks ended June 21, 1907. Estimated population, 31,600. Total number of deaths, 10. No deaths from contagious diseases reported.

St. Christopher.—Five weeks ended June 22, 1907. Estimated population, 49,289. Total number of deaths, 129, including 4 from tuberculosis.